

MSG BOF Meeting 9-19-12
Jim Harrington
CDFW Aquatic Bio-Assessment Lab
Cajun James
Sierra Pacific Industries













Why do we do bioassessments

How do we do bioassessment

How do we interpret bug data

How do we conduct bioassessments in California

Some examples and proposal for industrial timber lands

Why do we do bioassessments

How do we do bioassessment

How do we interpret bug data

How do we conduct bioassessments in California

Some examples and proposal for industrial timber lands

# California Department of Fish and Wildlife

Hot Creek Hatchery NPDES Permit





1993 in permit

Waited for response

1999-2004

2005 SI process

Continue monitoring

### 1993-95 Pilot Monitoring Study

### BIOLOGICAL CONDITION OF STUDY REACHES BASED ON MACRO-INVERTEBRATE FAUNAS

	Saara	
Score	_	Reach
Group 1 (COMPLEX)		
45		Upper Pudding Creek (Noyo Watershed)
43		Sweetwater Creek (Mokolumne Watershed)
39.5		Tiger Creek (Mokolumne Watershed)
37		Noyo River (Noyo Watershed)
35.5		Mill Creek (Mokolumne Watershed)
Group 2		
31.5		Doty Creek (Gualala Watershed)
29		Solinsky Crossing (Mokolumne Watershed)
Group 3 (SIMPLE)		
21		North Fork Gualala River (Gualala Watershed)
17.5		Lower Pudding Creek (Noyo Watershed)
13.5		Cottonwood Creek (Siskiyou Watershed)
12.5		Log Cabin Creek (Gualala Watershed)

Obviously, this ranking should be taken with a degree of caution. Without comparison to differences in habitat parameters, the biological significance of the groupings is not clear.

Subsampling 100 macro-invertebrates from a sample is recommended in the national RBPs (Plafkin et al 1989). The total number of macro-invertebrates in 11 samples examined ranged from 138 to 3950 with an average of 1140. Subsampling effected the metric values from 0 to 300% and the degree of difference was related to the difference in abundance (Appendix N, Table 7). In spring 1994, subsampling was increased to 300 to improve metric reliability. Metric values increased substantially (Appendix N, Table 2 and 3), but there was no difference in the coefficients of variation (Table 5 and 6). For 12 of the spring 1994 samples, there were less than 300 organisms. The lowest number was 134 but there was no noticeable effect to the metric value or variability.

A strong correlation could not be detected between bioassessment metric values and physical/habitat scores. This may be because the generalized visual evaluation was not quantitative or there was inconsistency in implementing the field procedures. The macro-invertebrate community appears to be more responsive to upstream influences than those in the immediate area. There were also inconsistencies between information indices generated for the streams visited in fall 1993 and spiring 1994.

The bioassessment metrics taxa richness, diversity index and EPT index seem to be reliable indicators of biological conditions. The biotic index had low variability which is indicative of a useful metric, but may not be a relevant metric for forested streams since it was developed for an indicator of organic pollution. Dominant taxa had the highest variability and its usefulness may be questionable.

The effectiveness of the physical/habitat evaluation procedure originally recommended by the EPA (Plafkin et al. 1989) was not determined in this study. Low-level ocular physical assessment

STREAM REACH	WATERSHE	D	SAMPLE I	D
RIFFLE #	DATE	TIME	WI	PCL #
WATER TEMP (°C) RIFFLE LENGTH (M)		HABITAT ASSESSMENT PAF (FILL IN AFTER FIELD DA (REFER TO HANDOUT FOR DEFIN		ELD DATA REVIEW)
TRANSECT INTE	ERSECT		1. BOTTOM SUBSTRA	ATE
#1 (m.cm)			2. INSTREAM COVER	_
#2 (м.см)			3. EMBEDDEDNESS	
#3 (m.cm)			4. VELOCITY / DEPTI	-
CREW MEMBERS		5. CHANNEL SHAPE		_
			6. POOL / RIFFLE RA	TIO
			7. WIDTH / DEPTH R	ATIO
			8. BANK VEGETATIO	N
			9. LOWER BANK STA	BILITY
			10. DISRUPTIVE PRE	SSURES
			11. ZONE OF INFLUE	NCE
COMMENTS				
CHAIN OF CUSTODY	RELEASED BY		RECEIVED BY DAT	DATE
COLLECTOR				
TRANSPORTER				
TRANSPORTER				
TRANSPORTER				

Sampled 11 sites in 3 watersheds
Purchased field and lab equipment
Tested field and lab methods
Modified US EPA methods for Cal needs
Was standard protocol until 2007

Ecological Integrity

Chemical Integrity

Physical Integrity

Biological Integrity

# Statutory Authority

- · Clean Water Act Section 101(a) Purpose:
  - "To restore and maintain the chemical, physical and biological integrity of the Nation's waters"





Biological integrity - the capability of the waterbody to support and maintain a balanced, integrated, adaptive community of organisms having a composition and diversity comparable to that of natural habitats of the region



Why do we do bioassessments

How do we do bioassessment

How do we interpret bug data

How do we conduct bioassessments in California

Some examples and proposal for industrial timber lands

# FRESHWATER BIOASSESSMENT - THE USE OF AQUATIC ORGANISM TO MEASURE AQUATIC HEALTH -

US EPA Recommendation:

Multiple Assemblages Fish, Invertebrates and Algae










# SWAMP Biossessment Procedures













Why do we do bioassessments

How do we do bioassessment

How do we interpret bug data

How do we conduct bioassessments in California

Some examples and proposal for industrial timber lands


### End Product - 600 subsampled BMIs from the sample reach



## Sensitive Organisms in Streams

**Dragonflies and Damselflies** 







**Stoneflies** 





**Caddisflies** 



Expected Response to Stress:



abundance & proportion

## Tolerant Organisms in Streams

Scucs



Leeches



Snails



Midges



**Expected Response to Stress:** 



# Types of BMI Metrics

Richness Measures

EPT Taxa

Composition Measures

Percent EPT Individuals

Tolerance/Intolerance Measures

Percent Sensitive EPT Taxa

Functional Feeding Groups

Percent Shredder Taxa

Total of 134

SAFIT Standard Taxonomic Effort I & II

## Algae Bioassessment: Supporting Documents







**Draft Technical Repor** 

2008

Incorporating Bioassessment Using Freshwater Algae into California's Surface Water Ambient Monitoring Program (SWAMP)

California's 
'Algae Plan'

March 2008









SWAMP Algae Field SOP

Standard Operating Procedure
for Collecting Stream Algae Sa
and Associated Physical Habits
and Chemical Data for Ambien
Bioassessments in California

#### July 2009

#### A. Elizabeth Fetscher

Southern California Coastal Water Research Project 3535 Harbor Blvd., Suite 110 Costa Mesa. CA 96262

#### Lilian Dune

San Diego Regional Water Quality Control Board State Water Resources Control Board 9174 Sky Park Court San Diego, CA 92123

#### Pete Ode

Aquatic Bioassessment Laboratory/Water Pollution Control Laboratory Department of Fish and game 2005 Nimbus Road Rancho Cordova, CA 95670

SWAMP
Algae
Field SOP
June 2009
(updated
May 2010)





Why do we do bioassessments


How do we do bioassessment


How do we interpret bug data


How do we conduct bioassessments in California

Some examples and proposal for industrial timber lands


# Targeted Sampling Site from Various Special Studies









### Biological condition of our nation's streams (USEPA 2006)



# 2007 CDFW Partnership with SWAMP Surface Water Ambient Monitoring Program



Water Boards




More than 2000 Random Sample Site through Combined Programs

2000 - 2003 US EPA


2004 - 2006 SWRCB Non-point Source Program


2007 - 2015 SWAMP




# **Reference Sites**


REGION	n
North Coast	75
Central Valley	1
Coastal Chaparral	57
Interior Chaparral	33
South Coast Mountains	85
South Coast Xeric	34
Western Sierra	131
Central Lahontan	114
Deserts + Modoc	27
TOTAL	586





Trout Creek
Tahoe Basin







Component	Obs	Expect
CSCI	0.65	1
O/E	0.49	1
0	7	14.1
MMI	0.81	1

Metric	Obs	Expect
% Coleoptera taxa	3	4
Diptera taxa	5	5.3
% EPT taxa	24	63
% Intolerant	5	36
% Non-insect	35	3
% Predator taxa	65	26
% Scraper taxa	0	10
% Shredders	0	10
Simpson's diversity	0.90	0.84
Tolerant taxa	6	5

# **Trout Creek Tahoe Basin**

Observed taxa	Missing taxa
Micrasema	Hydropsyche
Sweltsa	Diamesinae
Paraleptophlebia	Fallceon
Oligochaeta	Epeorus
Baetis	Rithrogena
Chironominae	Ameletus
Acari	Cinygmula
Orthocladiinae	Zapada
	Serratella
	Tanypodinae
	Rhyacophila
	Simulium
	Drunella





Technical Infrastructure

Regulatory Framework

Regulatory Applications

**SWAMP** 

**Standards** 

Regulatory Programs

Why do we do bioassessments

How do we do bioassessment

How do we interpret bug data


How do we conduct bioassessments in California


Some examples and proposal for industrial timber lands

# USFS Aquatic Management Indicator Species

- Lawsuit on EIS saying MIS not effective, No Data
- Decided to Change from Trout to BMIs
- 2009 first probabilistic sampling event (contractor issues)
- 2010 second probabilistic sampling event (contractor issues)
- 2011 and 2012 regroup contract with ABL
- 2013 ABL sampled 21 stream sites incorporated into PSA

### **2013 USFS Aquatic MIS Results**





### **Based on CSCI Scores**

USFS & CDFW combined indicate that 78 ± 6% of perennial stream miles on Sierra Nevada National Forests are in reference condition.



MSG BOF Meeting 9-19-12
Jim Harrington
CDFW Aquatic Bio-Assessment Lab
Cajun James
Sierra Pacific Industries















### King Fire Sampling Locations -Above, Within -Pilot Creek



### Ponderosa Fire Sampling Locations-Above, Within, Below



# Cooperative Pilot Project to Assess Stream Water and Habitat Quality of California Private Forest Lands April 2, 2015

### **Objective**

The intent of this project is to establish a collaborative monitoring framework for applying California's SWAMP ecological performance measures to evaluate water and habitat quality in streams on private forest lands. Direct collaborators include the State Water Resources Control Board, Department of Fish and Wildlife, Department of Forestry and Fire Protection, California Forestry Association, and private industrial forest owners.

#### **Background**

Over the past decade California has adopted ecological performance measures to evaluate resource management practices and support effective regulatory policies. Recently, wadeable streams have been the focus of a large multiagency effort to develop standardized technical and regulatory tools for measuring and assessing biological integrity as ecological performance measures. The State Water Board is currently developing guidance for the application of these tools that will become part of its Inland












## Ponderosa Fire – August 2012





And your Beetles

